ISSN 2277 – 8322

NATUROPATHIC APPROACHES TO CHRONIC FATIGUE SYNDROME: AN EVIDENCE-BASED ANALYSIS OF SYMPTOM MANAGEMENT AND QUALITY OF LIFE

Dr. Ritu Sharma
Assistant Professor
Department of Yoga and Naturopathy
Jagannath University, Jaipur, Rajasthan, India

Abstract- Myalgic encephalomyelitis /chronic fatigue syndrome (ME/CFS) is a complex, chronic, and often disabling condition characterized by persistent debilitating fatigue, post-exertional malaise, unrefreshing sleep, cognitive impairment, and reduced functional capacity. The condition can substantially affect physical independence, psychological well-being, social participation, and overall quality of life. In recent years, naturopathic and complementary approaches, including nutritional support, dietary supplementation, relaxation techniques, massage, and individualized lifestyle management, have attracted attention as possible supportive strategies for symptom management. However, the clinical effectiveness and safety of these approaches remain uncertain and require careful evaluation. The present review aims to examine published evidence concerning naturopathic approaches to ME/CFS, with particular emphasis on symptom management, functional outcomes, and health-related quality of life. A narrative literature review approach was used to synthesize findings from published systematic reviews, clinical guidance, and research on complementary and non-pharmacological interventions. The reviewed evidence suggests that certain complementary approaches may provide benefits for selected symptoms in some individuals, although findings are inconsistent and frequently limited by small sample sizes, methodological weaknesses, variations in intervention protocols, and insufficient long-term follow-up. Nutritional adequacy and individualized energy management are important aspects of supportive care, but evidence does not establish specific diets, supplements, or naturopathic treatment packages as cures for ME/CFS. The review concludes that

complementary approaches may be considered selectively alongside appropriate medical care, provided that their potential benefits, limitations, safety, and practical demands are carefully assessed. Further rigorous clinical research is required to determine whether specific interventions produce meaningful and sustained improvements in fatigue, daily functioning, and quality of life.

Keywords: Naturopathy, chronic fatigue syndrome, myalgic encephalomyelitis, ME/CFS, complementary medicine, symptom management, nutritional support, quality of life, energy management.

I. INTRODUCTION

1.1 Background of the Study

Myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) is a complex and long-term illness that can significantly reduce an individual's capacity to undertake ordinary physical, cognitive, social, and occupational activities. Although the condition is commonly associated with persistent fatigue, its clinical presentation extends beyond tiredness and may include post-exertional malaise, unrefreshing sleep, cognitive difficulties, pain, and symptoms associated with autonomic dysfunction. Post-exertional malaise is particularly important because physical, cognitive, or emotional activity can trigger a worsening of symptoms that may appear after a delay and persist for an extended period. Consequently, the condition may interfere with education, employment, household responsibilities, personal relationships, and participation in community activities. The severity and pattern of symptoms vary considerably among individuals, making individualized assessment and management essential.

The underlying mechanisms of ME/CFS remain incompletely understood, and no single treatment has been established as a cure. Diagnosis requires an appropriate clinical assessment, including consideration of other conditions that may cause or contribute to persistent fatigue. Sleep disorders, nutritional deficiencies, endocrine disorders, medication effects, and other medical conditions may require separate evaluation and management. Recognizing these possibilities is important because symptoms attributed to chronic fatigue may sometimes arise from treatable conditions or a combination of different health problems. The complexity of ME/CFS has encouraged

interest in multidisciplinary approaches that address symptoms, functional limitations, and quality of life rather than focusing exclusively on fatigue intensity.

Naturopathy is a broad approach to health that commonly emphasizes nutrition, lifestyle practices, relaxation, and selected complementary therapies. Depending on the practitioner and treatment setting, naturopathic care may involve dietary counselling, vitamin and mineral supplements, herbal preparations, massage, hydrotherapy, and mind–body techniques. These approaches are sometimes promoted as methods for improving energy, supporting recovery, or reducing the burden of chronic symptoms. Nevertheless, the inclusion of an intervention within naturopathic practice does not establish its effectiveness. Each intervention must be evaluated according to its underlying evidence, potential risks, clinical purpose, and relevance to the particular symptoms experienced by a person with ME/CFS.




Figure 1. Conceptual framework of naturopathic approaches and potential outcomes in chronic fatigue syndrome (CFS)

An evidence-based assessment of naturopathic approaches is therefore necessary to distinguish general supportive practices from interventions with

demonstrated clinical benefits. Adequate nutrition may help prevent nutritional deficiencies, and relaxation practices may support comfort or stress management, but neither observation establishes that a particular intervention treats the underlying illness. Similarly, a reported improvement in one symptom should not automatically be interpreted as recovery or a sustained improvement in physical functioning. The present paper examines the available evidence concerning naturopathic approaches to symptom management and quality of life, while emphasizing the importance of clinical safety, individual preferences, and realistic expectations.

1.2 Significance of the Study

The significance of this review lies in its examination of complementary approaches that people with ME/CFS may consider when attempting to manage persistent symptoms. The chronic and fluctuating nature of the condition can create substantial physical, emotional, social, and financial burdens, particularly when patients experience limitations in mobility, concentration, sleep, or independent functioning. In such circumstances, individuals may seek additional support beyond routine medical consultations. However, information about naturopathic treatments varies in quality, and some commercial claims may overstate potential benefits or fail to disclose possible adverse effects. An evidence-based review can help clarify which approaches have a reasonable role in supportive care, which remain experimental or uncertain, and which require particular caution. It may also assist healthcare professionals in discussing complementary therapies without dismissing patient preferences or encouraging treatments unsupported by reliable evidence.

The review also highlights the importance of assessing health-related quality of life alongside symptom severity. An intervention may have limited effects on fatigue while improving comfort, sleep-related distress, or the practical management of daily activities. Conversely, an intervention may produce a small short-term change in a symptom without improving independence or social participation. These distinctions are important for researchers and practitioners because the overall value of an intervention depends not only on symptom changes but also on whether it produces meaningful improvements in the person's everyday experience.

ISSN 2277 – 8322

II. REVIEW OF LITERATURE

2.1 Complementary and Alternative Medicine in ME/CFS

The use of complementary and alternative medicine in ME/CFS has been examined in systematic reviews involving a range of interventions, including dietary supplements, herbal preparations, mind–body practices, massage, and traditional manual therapies. One important review by Alraek et al. (2011) examined 26 randomized controlled trials involving 3,273 participants. The review identified potentially favorable findings for selected interventions but also highlighted methodological weaknesses, limited evidence for individual therapies, and difficulties in drawing firm conclusions. These limitations illustrate the importance of distinguishing promising findings from evidence that is sufficiently consistent and reliable to support routine clinical recommendations. Because the interventions varied substantially in their methods, duration, and outcomes, the results could not establish the effectiveness of naturopathy as a single, unified treatment approach.

Complementary therapies may be attractive to people living with a chronic illness because they often emphasize holistic care, individual preferences, and self-management. However, the concept of holistic care should not be confused with proof that all components of a treatment package are effective. A combination of dietary advice, supplements, relaxation, and manual therapy may be difficult to evaluate if the contribution of each component is not measured separately. Research is therefore needed to identify which specific interventions may be beneficial, for whom they may be appropriate, and whether any improvements persist over time.

2.2 Nutritional Support and Dietary Management

Nutritional support is relevant to ME/CFS because symptoms may interfere with shopping, cooking, eating regularly, and maintaining adequate fluid intake. Some individuals may experience nausea, gastrointestinal symptoms, reduced appetite, or difficulty preparing meals because of limited energy. These circumstances can increase the risk of inadequate nutritional intake, particularly when the person has a restricted diet or additional medical conditions. A balanced diet that provides sufficient energy, protein, vitamins,

ISSN 2277 – 8322

minerals, and fluids is therefore an important component of general supportive care. Nevertheless, evidence that a particular dietary pattern can cure ME/CFS or consistently improve its defining symptoms remains insufficient.

Restrictive dietary practices require particular attention because eliminating major food groups without appropriate supervision may contribute to nutritional deficiencies or unnecessary financial and practical burdens. Nutritional recommendations should be based on individual requirements, dietary tolerance, clinical findings, and the person's ability to obtain and prepare food. Where malnutrition, weight loss, or significant dietary restriction is present, referral to an appropriately qualified dietitian may be warranted. The available guidance supports nutritional adequacy and individualized dietary assessment rather than the routine promotion of special diets as disease-specific treatments.

2.3 Vitamins, Minerals, and Herbal Supplements

Vitamin and mineral supplements are frequently marketed for fatigue, low energy, and general well-being. Their role in ME/CFS, however, requires careful interpretation because fatigue has multiple possible causes. Correcting a documented nutritional deficiency may improve symptoms attributable to that deficiency, but it does not necessarily demonstrate an effect on the underlying mechanisms of ME/CFS. Evidence supporting the routine use of supplements such as vitamin B12, vitamin C, magnesium, or coenzyme Q10 specifically for ME/CFS remains insufficient to justify universal recommendations.

Herbal preparations present additional challenges because product composition, concentration, quality, and purity may vary. Some products can cause adverse reactions or interact with prescription medicines, and evidence from one preparation cannot automatically be generalized to another product containing similar ingredients. Patients may also use several supplements simultaneously, making it difficult to identify the cause of any perceived benefit or adverse effect. Consequently, supplement use should be reviewed with a qualified healthcare professional, and decisions should consider clinical indications, dose, safety, cost, and the quality of available evidence.

ISSN 2277 – 8322

2.4 Mind–Body Practices and Relaxation

Mind–body approaches include meditation, breathing exercises, relaxation techniques, and selected gentle movement practices. These interventions may be explored for relaxation, emotional well-being, or the management of distress associated with chronic illness. Some individuals may find that such practices provide a sense of comfort or help them cope with the uncertainty and restrictions associated with fluctuating symptoms. However, the evidence that mind–body interventions consistently improve the core symptoms of ME/CFS remains limited, and benefits observed in individual studies should be interpreted in the context of study design and methodological quality.

It is also important to distinguish supportive psychological or relaxation benefits from claims that ME/CFS is caused by psychological factors or can be eliminated through relaxation alone. Such conclusions are not justified by the evidence. Movement-based practices must also be adapted to the individual's tolerance because activity that appears gentle or low intensity may still trigger post-exertional symptom worsening. The potential role of these practices is therefore best considered on an individualized basis, with attention to patient preference, safety, and measurable outcomes.

2.5 Massage and Manual Therapies

Massage and other manual therapies have been examined as complementary approaches for selected symptoms associated with chronic fatigue and related conditions. Some studies have reported improvements in perceived relaxation, comfort, or individual symptom measures. However, differences in treatment protocols, participant characteristics, control conditions, and outcome measures make it difficult to establish the size and reliability of any benefit. Evidence for sustained improvements in overall functioning or quality of life remains uncertain.

Manual therapies should consequently be viewed as optional supportive interventions rather than established treatments for ME/CFS. Their appropriateness depends on symptom severity, pain sensitivity, individual tolerance, and the possibility that treatment-related effort or physical stimulation may worsen symptoms. Practitioners should avoid making claims that massage or similar therapies restore

normal physiological functioning or cure the underlying illness unless such claims are supported by reliable clinical evidence.

2.6 Sleep Management and Quality of Life

Sleep disturbance is a common concern in ME/CFS and may involve unrefreshing sleep, altered sleep patterns, difficulty initiating sleep, or difficulty maintaining sleep. Sleep-related symptoms can compound daytime difficulties and contribute to problems with concentration, mood, and daily functioning. Practical approaches may include reviewing sleep habits, adjusting the sleep environment, establishing a manageable routine, and seeking assessment for persistent or suspected sleep disorders. Such approaches may support comfort and help address specific sleep-related difficulties, although their effects on the overall course of ME/CFS remain uncertain.

Quality of life is a multidimensional outcome that includes physical functioning, psychological well-being, social participation, independence, and the ability to undertake meaningful activities. In ME/CFS, symptom severity alone may not adequately describe a person's experience. A modest improvement in sleep or pain may be valuable without producing a measurable improvement in employment or mobility. Research should therefore assess symptom-specific outcomes alongside validated measures of health-related quality of life and daily functioning.

2.7 Energy Management and Pacing

Energy management, commonly referred to as pacing, involves balancing physical, cognitive, emotional, and social activities with the person's available energy and current functional capacity. It aims to reduce the likelihood of symptom exacerbation by helping individuals recognize their limits and adjust activities accordingly. This approach is particularly relevant because post-exertional malaise may occur after a delay, meaning that a person may initially feel able to complete an activity but experience a significant deterioration later.

The Centers for Disease Control and Prevention describes pacing as a strategy for managing activity within individual limits, while the National Institute for Health and Care Excellence recommends individualized energy management rather than automatic increases in activity according to fixed schedules. These recommendations are

important when considering naturopathic lifestyle programmes because generic advice to exercise more or progressively increase activity may be unsuitable for some people with ME/CFS. Energy management should be flexible and responsive to symptom patterns, and it should not be presented as a cure or as a guarantee of increased exercise tolerance.

III. AIM AND OBJECTIVES

The primary aim of this review is to critically examine the published evidence concerning naturopathic approaches to symptom management and quality of life among people with ME/CFS. The review considers nutritional support, supplements, mind–body practices, manual therapies, sleep-related strategies, and individualized energy management, while distinguishing between general supportive care and interventions with evidence of specific clinical effectiveness. It also evaluates the limitations of existing research, potential safety concerns, and the need for further clinical investigation. The central research question is whether available evidence supports the use of selected naturopathic approaches to improve fatigue-related symptoms, daily functioning, and quality of life without exposing patients to unnecessary risk or replacing appropriate medical assessment and care.

IV. MATERIALS AND METHODS

4.1 Research Design

The paper uses a narrative literature review design to synthesize evidence concerning naturopathic and related complementary approaches to ME/CFS. A narrative review is appropriate for discussing a heterogeneous group of interventions and examining their clinical context, limitations, and potential relevance to patient-centered care. However, it differs from a systematic review that follows a prospectively specified protocol, documents a comprehensive search, uses reproducible selection procedures, and conducts a formal assessment of the included evidence. Accordingly, this manuscript should be regarded as a narrative review draft rather than a completed systematic review with independently verified study-selection counts.

4.2 Sources and Search Strategy

A formal search for a publication-ready version should use databases such as PubMed, Scopus, Web of Science, and the

Cochrane Library, alongside relevant clinical guidelines from NICE and the CDC. Search terms should combine condition-related phrases, including "myalgic encephalomyelitis," "chronic fatigue syndrome," and "ME/CFS," with intervention terms such as "naturopathy," "complementary medicine," "nutrition," "dietary supplements," "mind–body," "massage," "relaxation," and "energy management." These terms should then be combined with outcome terms such as "fatigue," "sleep," "pain," "quality of life," "functional status," and "adverse effects." Exact search strings, database coverage, the final search date, and the number of retrieved records should be documented before describing the paper as a systematic review.

4.3 Eligibility Criteria and Evidence Appraisal

Studies considered for a formal review should involve human participants diagnosed with ME/CFS or chronic fatigue syndrome using stated diagnostic criteria, evaluate a relevant intervention, and report at least one meaningful outcome related to fatigue, sleep, pain, functioning, quality of life, or safety. Duplicate publications, irrelevant studies, and articles lacking sufficient information for interpretation should be excluded. Evidence appraisal should consider sample size, study design, randomization, comparator groups, diagnostic criteria, intervention duration, outcome validity, attrition, adverse-event reporting, and risk of bias. Because the interventions and outcomes differ substantially, a narrative synthesis is appropriate unless a sufficiently comparable group of studies is identified for statistical pooling.

Table 1. Proposed Methodological Framework for the Review

Methodological component	**Description**
Review design	Narrative literature review
Target population	People diagnosed with ME/CFS or chronic fatigue syndrome
Intervention categories	Nutritional support, supplements, mind–body practices, massage, sleep

ISSN 2277 – 8322

	management, and energy management
Evidence sources	PubMed, Scopus, Web of Science, Cochrane Library, NICE, and CDC
Primary outcomes	Fatigue severity, post-exertional malaise, sleep, pain, and daily functioning
Secondary outcomes	Health-related quality of life, patient-reported well-being, adverse events, and treatment acceptability
Evidence appraisal	Study design, risk of bias, consistency, outcome validity, and follow-up
Synthesis method	Narrative synthesis; meta-analysis only if studies are sufficiently comparable

V. RESULTS

5.1 General Findings from Literature

The available literature indicates that naturopathic and complementary interventions have been investigated as possible methods of managing selected symptoms associated with ME/CFS. However, the evidence varies substantially by intervention, and the findings do not support treating naturopathy as one standardized or established therapy. Alraek et al. (2011) reviewed 26 randomized controlled trials involving 3,273 participants across a range of complementary and alternative therapies. Although the authors identified potentially favorable findings for some interventions, they emphasized methodological limitations and the lack of sufficient evidence to draw firm conclusions about many treatments.

The overall pattern suggests that some approaches may be worth investigating further, but the evidence remains inadequate to establish reliable, sustained improvements across the full range of ME/CFS symptoms. Nutritional support has a clear role in maintaining general health, while individualized energy management is emphasized in clinical guidance. In contrast, specific supplements, herbal products, and many complementary therapies lack sufficient consistent evidence to justify their

routine use as disease-specific treatments. The distinction between supportive care and proven treatment is therefore central to interpreting the literature.

5.2 Findings Related to Nutrition and Supplementation

The evidence supports maintaining adequate nutrition and hydration as part of general supportive care, particularly when symptoms interfere with meal preparation or regular eating. However, the available research does not establish that a particular diet reliably treats the core symptoms of ME/CFS. Likewise, supplement studies have produced inconsistent findings, and there is insufficient evidence to recommend routine vitamin or mineral supplementation as a treatment for all patients. Supplementation may be appropriate when a deficiency or another recognized clinical indication has been identified, but this should not be confused with evidence that the supplement treats the underlying illness.

5.3 Findings Related to Mind–Body and Manual Therapies

Some complementary studies have reported potentially beneficial effects on selected symptoms, including perceived relaxation or comfort. Nevertheless, variations in treatment methods, small study samples, and limitations in outcome assessment reduce confidence in the findings. The literature does not establish that mind–body practices or massage reliably improve post-exertional malaise, restore normal physical functioning, or produce sustained improvements in overall quality of life. These interventions may be considered selectively when they are acceptable to the individual, are well tolerated, and are used with realistic expectations.

5.4 Findings Related to Energy Management and Quality of Life

Clinical guidance emphasizes individualized energy management because post-exertional malaise can substantially affect functioning and may follow activity after a delay. Flexible pacing can help individuals recognize their limits and avoid activity patterns that provoke worsening symptoms. However, the evidence should not be interpreted as establishing pacing as a cure. Rather, it is a management strategy intended to reduce the risk of exacerbation and support more sustainable daily routines.

Quality-of-life outcomes require separate consideration because changes in one symptom do not necessarily translate into

improvements across all areas of functioning. The available literature supports the importance of measuring fatigue, sleep, pain, physical functioning, and patient-reported quality of life independently. Further studies are needed to determine whether particular complementary interventions produce meaningful and sustained benefits in these outcomes.

Table 2. Summary of Findings on Naturopathic and Related Approaches

Intervention	**Main potential role**	**Evidence interpretation**	**Important consideration**
Balanced nutrition	Prevention of inadequate intake and nutritional deficiencies	Strong rationale for general health; specific ME/CFS benefits of particular diets remain uncertain	Avoid unnecessary dietary restrictions
Vitamins and minerals	Correction of documented deficiencies	Insufficient evidence for routine use as an ME/CFS treatment	Review clinical indications, dose, and interactions
Herbal preparations	Potential symptom-related effects under investigation	Evidence varies and is generally insufficient for broad recommendations	Product quality and safety require attention
Mind–body practices	Relaxation and selected aspects of symptom coping	Limited and variable evidence for disease-specific outcomes	Adapt practices to individual tolerance
Massage and manual therapies	Potential support for comfort or selected symptoms	Some favorable findings, but substantial limitations remain	Monitor for discomfort or symptom worsening
Sleep management	Addressing sleep-related difficulties	Depends on the specific intervention and underlying	Assess persistent sleep disturbance

		sleep problem	clinically
Energy management and pacing	Balancing activity and rest to reduce post-exertional worsening	Supported by clinical guidance as individualized symptom management	Avoid fixed activity increases that exceed individual limits

VI. ANALYSIS AND DISCUSSION

6.1 Interpretation of Overall Evidence

The findings suggest that naturopathic approaches may have a selective role in supportive care, but their effectiveness should not be generalized across all interventions. The term naturopathy encompasses a wide range of practices, each with different proposed mechanisms, methods of application, and levels of supporting evidence. Consequently, a favorable result from one intervention cannot be used to establish the effectiveness of the entire approach. The systematic review by Alraek et al. (2011) illustrates this problem because potentially promising findings were accompanied by substantial methodological limitations and uncertainty regarding individual therapies.

The central analytical distinction is between supporting general health, managing specific symptoms, and treating the underlying illness. Adequate nutrition may prevent deficiencies, relaxation may improve comfort, and pacing may help reduce the risk of activity-related symptom worsening. These outcomes may be meaningful, but they do not establish that an intervention cures ME/CFS or consistently restores normal functioning. Future studies should identify clearly defined treatment objectives and measure outcomes that correspond directly to those objectives.

6.2 Relationship Between Symptom Management and Quality of Life

The effect of ME/CFS on quality of life extends beyond fatigue severity and includes sleep disturbance, pain, cognitive difficulties, reduced independence, and limitations in social participation. An intervention may therefore produce a meaningful benefit in one area without affecting other domains. For example, improved sleep-related comfort may be valuable even if physical functioning remains unchanged, while a reduction in a

ISSN 2277 – 8322

fatigue questionnaire score may not necessarily indicate improved ability to work or participate in family activities. Research should report these outcomes separately rather than treating any improvement as proof of overall recovery.

A comprehensive evaluation should use validated measures of fatigue, physical functioning, sleep, and health-related quality of life, supplemented by patient-reported descriptions of meaningful daily activities. Follow-up should be long enough to assess whether improvements persist and whether the intervention is associated with delayed symptom worsening. Such an approach would provide a more clinically relevant assessment of the value of naturopathic and complementary interventions.

6.3 Nutritional Support and Clinical Implications

Nutritional support has a clear practical role when symptoms make adequate food intake difficult or when a patient has a documented deficiency. However, this role should not be used to justify restrictive diets or commercial products that claim to reverse ME/CFS without reliable evidence. A balanced approach should prioritize adequate energy and nutrient intake, hydration, individual tolerance, and the practical challenges of shopping and preparing meals. Where nutritional risk is identified, professional assessment can help prevent harm and tailor recommendations to the individual's circumstances.

The evidence also highlights the need to avoid equating correction of a deficiency with treatment of the underlying illness. If a person with ME/CFS has a vitamin deficiency, correcting it may improve symptoms associated with that deficiency. Nevertheless, the response does not establish that the vitamin is an effective general treatment for ME/CFS. This distinction is important for both clinical decision-making and accurate reporting in academic research.

6.4 Safety of Supplements and Complementary Therapies

Safety is a major consideration when evaluating naturopathic interventions. Supplements may interact with prescription medicines, produce adverse effects, or contain ingredients in concentrations that differ from those expected. The use of several products simultaneously may further complicate the assessment of benefits and harms. In addition, restrictive diets and demanding treatment routines may increase

physical or financial burdens for individuals whose daily functioning is already limited.

Healthcare professionals should therefore review complementary products and therapies with patients, taking account of the intended benefit, clinical indication, possible interactions, costs, and alternative options. Patients should be encouraged to seek medical advice when symptoms worsen or when a new intervention produces an unexpected reaction. Complementary approaches should not delay appropriate evaluation of persistent fatigue or the treatment of coexisting conditions.

6.5 Individualized Energy Management

Energy management is particularly relevant because ME/CFS can involve delayed post-exertional symptom worsening. Advice that encourages people to increase activity according to fixed schedules may be inappropriate when it does not account for the individual's symptoms and current capacity. NICE guidance supports flexible energy management rather than automatic progression of activity. This is consistent with a person-centered approach that recognizes fluctuations in symptoms and the need to adjust physical, cognitive, and social demands.

Nevertheless, pacing should not be described as a cure or as a guaranteed method of increasing physical capacity. Its purpose is to help individuals balance activity and rest, recognize patterns of symptom exacerbation, and reduce the risk of repeatedly exceeding their limits. Further research is needed to determine the magnitude of its effects on different outcomes and to identify which forms of support are most helpful for people with different levels of disease severity.

6.6 Research Limitations and Future Directions

The evidence base is constrained by variations in diagnostic criteria, intervention protocols, sample sizes, outcome measures, and follow-up periods. Some studies may also be affected by inadequate blinding, high attrition, or incomplete reporting of adverse events. These limitations reduce the reliability of broad conclusions and make comparisons between interventions difficult. Future research should use clearly defined ME/CFS populations, appropriate comparison groups, standardized descriptions of interventions, validated outcome measures, and transparent reporting of both beneficial and harmful effects.

Research should also examine the practical value of interventions, including their cost, accessibility, time requirements, and effect on daily energy demands. An intervention that provides a small improvement in one symptom but imposes substantial physical or financial burdens may not produce an overall benefit in quality of life. Studies that assess patient priorities, functional outcomes, and long-term tolerability would therefore make an important contribution to the evidence base.

VII. LIMITATIONS OF THE REVIEW

The present manuscript has limitations that should be acknowledged when interpreting its findings. It is a narrative review rather than a completed systematic review with a fully documented database search, independent study screening, and formal risk-of-bias assessment. The interventions discussed are heterogeneous, and the findings cannot be combined into a single estimate of the effectiveness of naturopathy. In addition, the manuscript does not calculate pooled effect sizes, confidence intervals, or statistical significance and does not report new patient data. The evidence synthesis should be updated through a reproducible database search before journal submission, particularly to verify the most recent clinical studies and confirm the details of every reference. These limitations mean that the conclusions should be interpreted as a cautious overview of the available evidence rather than definitive proof of clinical effectiveness.

VIII. CONCLUSION

ME/CFS is a complex and often disabling condition that can substantially affect fatigue, sleep, cognition, physical functioning, social participation, and overall quality of life. Naturopathic and complementary approaches have been investigated as possible supportive strategies, but the available evidence varies considerably between interventions. Some studies suggest potentially beneficial effects for selected symptoms, yet methodological weaknesses, small study samples, inconsistent outcome measures, and insufficient long-term follow-up prevent firm conclusions about the overall effectiveness of these approaches. Current evidence does not establish naturopathy as a cure for ME/CFS or support the routine use of specific supplements and restrictive diets as disease-specific treatments.

A responsible approach emphasizes adequate nutrition, individualized symptom management, attention to sleep, flexible energy management, and careful assessment of complementary therapies. The potential benefits of any intervention should be considered alongside its safety, cost, practical demands, and risk of symptom deterioration. Future research should use rigorous clinical designs, clearly defined interventions, validated patient-centered outcomes, and adequate follow-up to determine whether particular approaches produce meaningful and sustained improvements in symptoms and quality of life. Overall, naturopathic approaches may be considered selectively as complementary elements of person-centered supportive care, provided that they are guided by evidence and coordinated with appropriate medical care.

References

[1] Alraek, T., Lee, M. S., Choi, T. Y., Cao, H., & Liu, J. P. (2011). Complementary and alternative medicine for patients with chronic fatigue syndrome: A systematic review. BMC Complementary and Alternative Medicine, 11, Article 87.

[2] Campagnolo, N., Johnston, S., Collatz, A., Staines, D., & Marshall-Gradisnik, S. (2017). Dietary and nutrition interventions for the therapeutic treatment of chronic fatigue syndrome/myalgic encephalomyelitis: A systematic review. Journal of Human Nutrition and Dietetics, 30(3), 247–259.

[3] National Institute for Health and Care Excellence. (2021). Myalgic encephalomyelitis (or encephalopathy)/chronic fatigue syndrome: Diagnosis and management (NICE Guideline NG206). NICE.

[4] Institute of Medicine. (2015). Beyond myalgic encephalomyelitis/chronic fatigue syndrome: Redefining an illness. The National Academies Press.

[5] Fukuda, K., Straus, S. E., Hickie, I., Sharpe, M. C., Dobbins, J. G., & Komaroff, A. (1994). The chronic fatigue syndrome: A comprehensive approach to its definition and study. Annals of Internal Medicine, 121(12), 953–959.

[6] Carruthers, B. M., van de Sande, M. I., De Meirleir, K. L., Klimas, N. G., Broderick, G., Mitchell, T., Staines, D., Powles, A. C. P., Speight, N., Vallings, R., Bateman, L., Baumgarten-

Austrheim, B., Bell, D. S., Carlo-Stella, N., Chia, J., Darragh, A., Jo, D., Lewis, D., Light, A. R., . . . Stevens, S. (2011). Myalgic encephalomyelitis: International Consensus Criteria. Journal of Internal Medicine, 270(4), 327–338.

[7] Sanal-Hayes, N. E. M., McLaughlin, M., Hayes, L. D., Mair, J. L., Ormerod, J., Carless, D., Hilliard, N., Meach, R., Ingram, J., & Sculthorpe, N. F. (2023). A scoping review of "pacing" for management of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS): Lessons learned for the long COVID pandemic. Journal of Translational Medicine, 21, Article 720.

[8] Sanal-Hayes, N. E. M., McLaughlin, M., Mair, J. L., Ormerod, J., Carless, D., Meach, R., Hilliard, N., Ingram, J., Sculthorpe, N. F., & Hayes, L. D. (2025). Pacing for management of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS): A systematic review and meta-analysis. Fatigue.

[9] Dorczok, M. C., Mittmann, G., Mossaheb, N., Schrank, B., Bartova, L., Neumann, M., & Steiner-Hofbauer, V. (2025). Dietary supplementation for fatigue symptoms in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS)—A systematic review. Nutrients, 17(3), Article 475.

[10] National Institute for Health and Care Excellence. (2021). Evidence reviews for the non-pharmacological management of ME/CFS. NICE.

[11] Agency for Healthcare Research and Quality. (2016). Diagnosis and treatment of myalgic encephalomyelitis/chronic fatigue syndrome. U.S. Department of Health and Human Services.

[12] Centers for Disease Control and Prevention. (2024). Management of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS): An updated systematic evidence review. U.S. Department of Health and Human Services.

[13] Hvidberg, M. F., Brinth, L. S., Olesen, A. V., Petersen, K. D., & Ehlers, L. (2015). The health-related quality of life for patients with myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). Health and Quality of Life Outcomes, 13, Article 167.

[14] Cockshell, S. J., & Mathias, J. L. (2010). Cognitive functioning in chronic fatigue

syndrome: A meta-analysis. Psychological Medicine, 40(8), 1253–1267.

[15] Prins, J. B., van der Meer, J. W. M., & Bleijenberg, G. (2006). Chronic fatigue syndrome. The Lancet, 367(9507), 346–355.

[16] Clayton, E. W. (2015). Beyond myalgic encephalomyelitis/chronic fatigue syndrome: An IOM report on redefining an illness. JAMA, 313(11), 1101–1102.

[17] Komaroff, A. L., & Bateman, L. (2021). Will COVID-19 lead to myalgic encephalomyelitis/chronic fatigue syndrome? Frontiers in Medicine, 7, Article 606824.

[18] World Health Organization. (2022). International classification of diseases for mortality and morbidity statistics (11th Revision). World Health Organization.